

Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance

*Shaltout, F. A.; **El-Diasty, E.M. and ***Mohamed, M. S.

*Faculty of Vet. Med., Moshtohr, Benha University

** Animal Health Research Institute Dokki, Giza

***Egypt food factories

Abstract

A total of 100 random samples of chicken meat products; 25 of chicken fillet, chicken luncheon, chicken pane and chicken minced meat; were collected and subjected to mycological evaluation. The mean total mould counts were $1.9 \times 10^2 \pm 8.2 \times 10^1$, $3.3 \times 10^2 \pm 2.0 \times 10^2$, $2.8 \times 10^2 \pm 1.4 \times 10^2$ and $1.9 \times 10^2 \pm 3.4 \times 10^1$ CFU / g for chicken fillet, chicken luncheon, chicken pane and chicken minced meat respectively. Respective yeast counts were $5.7 \times 10^2 \pm 3.7 \times 10^2$, $3.3 \times 10^2 \pm 2.0 \times 10^2$, $2.1 \times 10^2 \pm 1.1 \times 10^2$ and $4.3 \times 10^3 \pm 9.8 \times 10^2$ CUF/ g. The predominant mould genera isolated from chicken fillet, chicken luncheon, chicken pane and chicken minced meat were *Aspergillus* followed by *Penicillium* then *Geotrichum*, *Fusarium*; *Cladosporium*; *Mucor*; *Eupenicillium*; *Scopulariopsis* and *Acremonium*. The predominant species of yeasts isolated from chicken fillet, chicken luncheon, chicken pane and chicken minced meat were *Candida* spp. (32.5%) followed by *Rhodotorula* spp. (22.1%), *Saccharomyces* spp. (18.2%), *Torulopsis* spp. (15.5%) and *Cryptococcus* spp. (11.7%). The isolated moulds and yeasts were evaluated for proteolytic and lipolytic activities on skim milk agar and Tributyrin agar. The economic and public health significance of isolated moulds and yeasts as well as the sanitary precautions were discussed.

Key words:

Chicken meat products, mould, yeast, proteolytic, lipolytic

Introduction

Chicken and chicken products provide a source of animal protein of high biological value for consumers at all ages, where they contain all the essential amino acids required for growth with high proportion of unsaturated fatty acids and low cholesterol contents. Moreover, poultry meat is good source of different types of vitamins as niacin, riboflavin, thiamine and ascorbic acid as well as sodium, calcium, iron, phosphorus, sulphur and iodine.

In processing plants, contamination of poultry meat products can occur throughout ideal processing, packaging and storage until the product is sufficiently cooked and consumed. Heavy microbial loads enter the processing operations

with the living birds and these microorganisms can be disseminated throughout the plant during processing. Diseases can also result when these products are not properly cooked and post-processing contaminated (Zhang *et al.*, 2001).

Fungi are of considerably less importance in poultry spoilage except when antibiotics are employed to suppress bacterial growth. When antibiotics are employed, however, moulds become the primary agents of spoilage. The genera of *Candida*, *Rhodotorula*, *Debaryomyces*, and *Yarrowia* are the most important yeasts found on poultry meat (Jay, 1979).

However, moulds and yeasts are of great importance in spoilage of poultry meat products

resulting in different changes in flavour, colour, texture and odour and also these fungi responsible for major portion of food deterioration especially in poor developing countries due to lack of hygienic measure and due to the use of contaminated additives and spices which considered a major important sources of mould contamination (Abd El-Rahman, 1987).

Therefore, the present study was planned out to throw a light on the total mould and yeast counts of chicken products (pane, minced meat and luncheon), and determination of lipolytic and proteolytic activities of the existing moulds and yeast.

Materials and Methods

Collection of samples

A total of 100 random samples of processed chicken meat products (25 samples of each chicken fillet, pane, minced meat and luncheon) were collected from shops and supermarkets. These samples were obtained and preserved in an ice box, then transferred to the

laboratory under complete aseptic conditions and examined as rapidly as possible.

Fungal isolation and identification

Total fungal count was carried out according to the techniques recommended by ISO (217-1-2:2008). Fungi were isolated and identified according to macroscopic and microscopic characteristics as described by Pitt and Hocking (2009). Identifications of the yeast isolates were performed according to Kriger Van Rij (1984) and Deak, (2008).

Evaluation of lipolytic and proteolytic activities of existing moulds and yeasts isolated from examined samples.

Lipolytic activity was determined by using Tributyrin agar medium according to the technique recommended by Koburger and Jacger (1987). Proteolytic activity was determined by using a casein substrate as described by O'reilly and Day (1983).

Results

Table (1). Incidence of moulds and yeasts in examined chicken meat products (N= 25).

Examined samples	Chicken Fillet		Chicken Luncheon		Chicken Panne		Chicken Minced meat	
	No. of positive samples	%	No. of positive samples	%	No. of positive samples	%	No. of positive samples	%
Mould	15	60	16	64	25	100	20	80
yeast	21	84	20	80	17	68	25	100

Table (2). Statistical analytical results of total mould count/g of examined chicken meat products samples.

Examined samples	Minimum	Maximum	Mean $\pm$ SE.
Chicken Fillet	10	2×10^3	$1.9 \times 10^2 \pm 8.2 \times 10^1$
Chicken luncheon	20	4×10^3	$3.3 \times 10^2 \pm 2.0 \times 10^2$
Chicken Pane	20	3.5×10^3	$2.8 \times 10^2 \pm 1.4 \times 10^2$
Chicken Minced meat	20	8.1×10^3	$1.9 \times 10^2 \pm 3.4 \times 10^1$




Figure (1). Mean mould counts / g of examined chicken meat product samples.

Table (3). Statistical analytical results of total yeasts count/g of examined chicken meat products samples. (N=25)

Examined samples	Minimum	Maximum	Mean $\pm$ SE.
Chicken Fillet	20	2.9×10^3	$5.7 \times 10^2 \pm 3.7 \times 10^2$
Chicken luncheon	50	2.9×10^3	$3.3 \times 10^2 \pm 2.0 \times 10^2$
Chicken Pane	<10	2.6×10^3	$2.1 \times 10^2 \pm 1.1 \times 10^2$
Chicken Minced meat	1.2×10^2	1.7×10^4	$4.3 \times 10^3 \pm 9.8 \times 10^2$




Figure (2). Mean yeast counts / g of examine d chicken meat product samples




Figure (3). Frequency percentage of moulds species isolated from examined samples of chicken meat products.

Table (4). Incidence of *Aspergillus* species isolated from examined samples of different chicken meat products (N=55).

Product Aspergillus species	Chicken Fillet		Chicken Luncheon		Chicken Pane		Chicken Minced meat	
	No.	%	No.	%	No.	%	No.	%
<i>A. flavus</i>	3	5.5	5	9.1	7	12.8	6	10.9
<i>A. niger</i>	5	9.1	6	10.9	8	14.5	6	10.9
<i>A. ochraceus</i>	-	-	-	-	5	9.1	-	-
<i>A. terreus</i>	-	-	-	-	2	3.6	-	-
<i>A. clavatus</i>	-	-	-	-	1	1.8	-	-
<i>A. candidas</i>	1	1.8	-	-	-	-	-	-

N= Number of isolated *Aspergillus* species




Figure (4): Frequency percentage of *Aspergillus* species isolated from examined chicken meat product (according to number of isolated *Aspergillus* species)




Figure (5): Frequency percentage of Penicillium species isolated from examined chicken meat product (according to number of isolated Penicillium species)

Table (5): Incidence of yeast species isolated from examined samples of chicken meat products (N=79).

Products yeast species	Chicken Fillet		Chicken Luncheon		Chicken Pane		Chicken Minced meat	
	No.	%	No.	%	No.	%	No.	%
<i>Candida</i> spp.								
<i>C. albicans</i>	-	-	-	-	-	-	1	1.3
<i>C. famata</i>	-	-	1	1.3	1	1.3	1	1.3
<i>C. krusei</i>	3	3.8	1	1.3	1	1.3	1	1.3
<i>C. lusitaniae</i>	-	-	1	1.3	-	-	1	1.3
<i>C. parapsilosis</i>	2	2.6	1	1.3	1	1.3	2	2.6
<i>C. pelliculosa</i>	1	1.3	1	1.3	-	-	1	1.3
<i>C. tropicalis</i>	-	-	1	1.3	1	2.6	1	1.3
<i>Cryptococcus</i> spp.	-	-	-	-	1	2.6	3	3.8
<i>Cry. albidus</i>	-	-	1	1.3	1	1.3	-	-
<i>Cry. species</i>	-	1.3	2	2.6	1	1.3	3	3.8
<i>Saccharomyces</i> spp.								
<i>S. cerevisiae</i>	1	1.3	-	-	1	1.3	2	2.6
<i>S. species</i>	2	2.6	5	6.4	1	1.3	2	2.6
<i>Rhodotorula</i> spp.								
<i>Rho. glutinis</i>	1	1.3	2	2.6	2	2.6	2	2.6
<i>Rho. species</i>	3	3.8	2	2.6	1	1.3	4	5.1
<i>Torulopsis</i> spp.	2	2.6	3	3.8	2	2.6	5	6.4


N= Number of isolated yeast species

Table (6). Proteolytic and lipolytic activity of some isolated mould.

Mould species	Proteolytic activity	Lipolytic activity
<i>A. flavus</i>	+ve	+ve
<i>A. candidas</i>	+ve	+ve
<i>A. clavatus</i>	-ve	-ve
<i>A. ochraceus</i>	+ve	+ve
<i>A. niger</i>	+ve	+ve
<i>A. terreus</i>	+ve	+ve
<i>P. brevicompactum</i>	-ve	+ve
<i>P. citreonigrum</i>	+ve	+ve
<i>P. chrysogenum</i>	+ve	+ve
<i>P. corylophilum</i>	+ve	+ve
<i>P. decumbens</i>	-ve	+ve
<i>P. griseofulvum</i>	+ve	+ve
<i>P. purpurogenum</i>	+ve	+ve
<i>P. simplicissimum</i>	-ve	+ve
<i>P. thomii</i>	+ve	+ve

Table (7). Proteolytic and lipolytic activity of some isolated yeasts.

Yeast species	Proteolytic activity	Lipolytic activity
<i>C. albicans</i>	-ve	+ve
<i>C. famata</i>	-ve	+ve
<i>C. krusei</i>	+ve	+ve
<i>C. lusitaniae</i>	+ve	+ve
<i>C. parapsilosis</i>	+ve	+ve
<i>C. pelliculosa</i>	+ve	+ve
<i>C. tropicalis</i>	+ve	+ve
<i>Cry. albidus</i>	-ve	+ve
<i>S. cerevisiae</i>	-ve	+ve
<i>Rho. glutinis</i>	-ve	+ve

**Figure (6).** Show proteolytic and lipolytic activity of isolated moulds and yeasts, (A) Proteolytic activity of *Penicillium purpurogenum*, (B) Lipolytic activity of *Penicillium griseofulvum*, (C) Lipolytic activity of *A. flavus* and (D) Lipolytic activity of *C. krusei*

Discussion

Moulds only compete with bacteria on meat when storage temperatures are lowered to 0°C or below, or when the meat surface dries to an a_w that enables fungi to compete (Pitt and Hocking, 2009).

Several authors cited several reports of bacteria and fungi growing on meats at -5°C, with yeasts and moulds predominating as temperatures were further lowered, to a limit at about -12°C. Schmidt-Lorenz and Gutschmidt (1969) reported that moulds and yeasts grew on chickens stored at -7.5 and -10 ± 0.2°C for 1 year.

The result achieved in table (1) revealed that the incidence of moulds in the examined chicken meat product samples were 15(60%), 16 (64%), 25(100%) and 20(80%) for chicken filet, chicken luncheon, chicken pane and chicken minced meat respectively. While, the incidence of yeasts in the examined chicken meat product samples were 21(84%), 20 (80%), 17(68%) and 25(100%) for chicken filet, chicken luncheon, chicken pane and chicken minced meat respectively. The results obtained for chicken filet, chicken luncheon, chicken pane and chicken minced meat are similar to that recorded by many investigators (Abdel-Rahman *et al.*, 1985, Edris 1986 and Rwaih-Wadee, 2010) who stated that the incidence of moulds in the examined chicken meat product samples were 13 (86.6%), 14 (93.33%), 13 (86.6%) of chicken luncheon, chicken pane and chicken minced meat respectively.

From the results achieved in table (2) and figure (1) it is evident that the mean mould count in chicken filet was $1.9 \times 10^2 \pm 8.2 \times 10^1$ cfu/g, while chicken luncheon was $3.3 \times 10^2 \pm 2.0 \times 10^2$ cfu/g. Concerning the examined samples of chicken pane and chicken minced meat, it is evident from the results presented in figure (1) that the mean values of mould count $2.8 \times 10^2 \pm 1.4 \times 10^2$ and $1.9 \times 10^2 \pm 3.4 \times 10^1$ cfu / g, respectively.

The obtained results declared that the examined chicken luncheon samples had the highest

mould count followed by chicken pane samples, while the chicken minced meat had the lowest count.

The obtained results were higher than those reported by Rwaih-wadee (2010) who found that the mean value of total fungal counts in examined samples of chicken luncheon, chicken pane and chicken minced meat were 58.86 ± 14.44 , 44.60 ± 13.28 and 53.33 ± 21.25 cfu/g. While higher figures were revealed by Shaltout (1996), El-Deeb *et al.*, (2011), Shawish (2011), Sharaf and Sabra (2012), and Yousif *et al.*, (2013), El-diahy *et al.*, (2013) who reported that the mean values of the total mould count of the chicken meat slice, chicken luncheon and chicken minced meat were 2.51, 3.71 and 4.12 log CFU/g, respectively.

Regarding the results recorded in table (3) and figure (2) it is obvious that the mean yeast count of examined samples in chicken filet, chicken luncheon, chicken pane and chicken minced meat were $5.7 \times 10^2 \pm 3.7 \times 10^2$, $3.3 \times 10^2 \pm 2.0 \times 10^2$, $2.1 \times 10^2 \pm 1.1 \times 10^2$ and $4.3 \times 10^3 \pm 9.8 \times 10^2$ CUF/ g, respectively. The minimum counts were of 20, 50, <10 and 1.2×10^2 , respectively. While, the maximum count were 2.9×10^3 , 2.9×10^3 , 2.6×10^3 and 1.7×10^4 , respectively. Higher counts were obtained by Brr and Mahmoud (2005), Hassan (2007), Ouf *et al.*, (2010), El-Deeb *et al.*, (2011) and El-diahy *et al.*, (2013). Comparatively most of the previous researches had higher counts due to the differences between the chicken processing plant at this time (sanitation program – computerized machines – educated workers) and manufacturing in the past.

Chicken Luncheon and Chicken pane is exposed to heat treatment during its processing and any microbial activities may be recorded as a result of post processing contamination, unhygienic storage as well as long storage life which may lead to spoilage. Also, spices have been used in many meat industries and this may add more contamination to meat products. Having been dried material from plant origin, spices are commonly heavily contaminated

with xerophilic storage moulds. The most frequent fungal contaminants of spices are species from the genera *Aspergillus* and *Penicillium* (Kocić-Tanackov *et al.*, 2007).

The results given in figure (3) revealed that the chicken meat products samples were contaminated with many fungal genera and species. The predominant mould genera isolated from chicken fillet, chicken luncheon, chicken pane and chicken minced meat were *Aspergillus* 9 (36%), 11(44%), 23 (92%) and 12(48%) respectively, followed by *Penicillium* 3(12%), 9 (36%), 16 (64%), and 4(16%) then *Geotrichum* 10 (40%), 5(20%), 6(24%) and 7(28%); *Fusarium* 5 (20%), 2(8%), 3(12%) and 2(8%); *Cladosporium* 5 (20%), 0 (0%), 0 (0%) and 5 (20%); *Mucor* 4 (16%), 0 (0%), 0 (0%) and 5 (20%); *Eupenicillium* 0 (0%), 3(12%), 2(8%) and 1(4%); *Scopulariopsis* 0 (0%), 1(4%), 2 (8%) and 0 (0%); *Acremonium* 0 (0%), 1 (4%), 0 (0%) and 2(8%) Finally *Talaromyces* isolated only from Chicken Pane 1(4%). Nearly similar strains were recorded in different poultry meat products by Abdel-Rahman *et al.*, (1985), Edris (1986), El-Khateib and Abd El-Rahman (1989), Edris *et al.*, (1992), El-Gazzar (1995), Shaltout (1996), Shaltout (2002), Bkheet *et al.*, (2007), Hussein (2008), Shawish (2011) and El-Diasty *et al.*, (2013).

The present data in table (4) and figure (4) shows the number and percentages of *Aspergillus* isolated from examined samples, *A. niger* was 5 (9.1%), 6 (13.04%), 8 (17.4%) and 6 (13.04%) from Chicken Fillet, chicken luncheon, chicken pane and chicken minced meat samples respectively, followed by *A. flavus* 5 (5.5%), 5 (10.9%), 7 (15.21%) and 6(13.04%) respectively, *A. ochraceus*, *A. terreus* and *A. clavatus* detected in chicken pane only 5 (10.9%), 2(4.34%) and 1 (2.2 %) respectively. *A. candidas* 1 (1.8%) isolated only from Chicken Fillet.

The obtained results declared that the chicken pane and chicken minced meat samples had the highest number of *Aspergillus* spp. 23 and 12 isolates, respectively, followed by chicken luncheon (11 isolates). *Aspergillus* is a ubiquitous soil-dwelling fungus (Wong *et al.*, 2008).

Identifications of *Penicillium* spp. isolated from the examined chicken meat products in figure (5) revealed that *P. citreonigrum*, *P. simplicissimum* were the most predominant species isolated from the chicken luncheon, chicken pane and chicken minced meat samples, 1(3.3 %), 5(16.7%) and 1(3.3 %), respectively, and 4(13.3%), 4(13.3%) and 1(3.3 %), respectively. *Penicillium corylophilum* 1 (3.0%), 2(6.7%), 1(3.3 %) and 1(3.3 %), respectively in chicken fillet chicken, luncheon, chicken pane and chicken minced meat samples. *Penicillium thomii* and *P. decumbens* were 3(10%) and 1(3.3 %), respectively, and 1 (3.3 %) and 1(3.3%), respectively in chicken pane and chicken minced meat samples only. In chicken luncheon *P. griseofulvum* and *P. brevicompactum* 2(6.7%) and 1(3.3 %), respectively, *P. chrysogenum* 1(3.3 %) in chicken pane and *P. purpurogenum* were isolated from chicken pane 1(3.3 %).

Some fungi may produce secondary metabolites known as mycotoxins. Mycotoxins are potent animal carcinogens and have been classified by the International Agency for Research in Cancer (IARC, 1993) as human carcinogens or potential (probable and possible) human carcinogens.

Regarding the results tabulated in table (5) shows the incidence of yeast genera isolated from the examined samples of chicken fillet, chicken luncheon, chicken pane and chicken minced meat *Candida* spp. 25 (32.5%) was the predominant yeast species isolated from such samples followed by *Rhodotorula* spp. 17 (22.1%), *Saccharomyces* spp. 14 (18.2%), *Torulopsis* spp. 12 (15.5%) and *Cryptococcus* spp. 9 (11.7%).

Nearly similar results were recorded by Edris (1992), Hegazi *et al.*, (1992), Shaltout (1996), Viljoena *et al.*, (1998), Shaltout (2002), Brr and Mahmoud (2005), Deák (2008), Ouf *et al.*, (2010), Shawish (2011), Seham- Ismail (2013), El-Diasty *et al.*, (2013) and Samaha (2013).

Moulds and yeasts are capable of hydrolyzing a wide range of proteinaceous materials (Koburger, 1972). In this respect, the prote-

olytic activity of isolated moulds from examined chicken meat products was evaluated using litmus milk medium (Table, 6 and 7). The obtained results showed that *A. flavus*, *A. niger*, *A. ochraceus*, *A. terreus*, *A. candidus*, *P. corylophilum*, *P. griseofulvum*, *P. Citreonigrum*, *P. chrysogenum*, *P. simplicissimum*, *P. purpurogenum*, *P. thomii*, were having proteolytic activity with clear zone of casein hydrolysis. While, all *Candida* species had proteolytic activity except *C. albicans*, also *C. krusei*, *Cry. albidus*, *S. cerevisiae* and *Rho. glutinis*.

On the other hand the lipolytic activity revealed that nearly all the isolated fungi and species of *C. albicans*, *C. krusei*, *C. parapsilosis*, *C. pelliculosa*, *C. lusitanae*, *C. tropical*, *C. famata*, *Cry. albidus*, *S. cerevisiae* and *Rho. glutinis* were found to be lipolytic on the tributyrin agar media. These results nearly similar to those obtained by Abd El-Rahman and Saad (1989), Welthagen and Viljoen (1999), Abd El-Aziz (1999), Farag (2000), Nasser (2002), Shaltout (2002), Subash, *et al.*, (2005), Soliman and Shalaby (2008), Korashy and Wahbba (2008), El-Diasty and Salem (2009) and Ouf *et al.*, (2010).

In conclusion, the obtained results showed high contamination of chicken meat products with different types of yeasts and moulds which constitute a public health hazard. Obviously, it is important to prevent mould growth to avoid toxin production through preventing the natural contamination of raw materials. Storage of food under conditions which prevent mould growth and strict hygienic measures and regulations should be imposed during processing, packing and transportation.

References

- Abd El-Aziz, N.M. (1999). Mycological status of meat and some meat products. Ph.D. Thesis. Meat hygiene. Fac. Vet. Med. Ass. Univ., Egypt.
- Abd El-Rahman, H. (1987). "Mycological studies on some selected species with special references to aflatoxin producing *Aspergillus flavus* species". Assuit Vet. Med. J., 19 (37): 93-100.
- Abdel-Rahman, H.; Darwish, A. and Hamdy, M. (1985). Mould affection of a meat cold store. Assuit Vet. Med. J., 14 (27): 131-134.
- Abd El-Rahman, H.A. and Saad, S.M. (1989). Studies on proteolytic activity of some prevalent mold species in relation to meat hygiene. Zagazig Vet. Med. J., 17(2): 228- 236.
- Bkheet, A.A.; Rezk, M.S. and Mousa, M.M. (2007). study on the microbiological content of local manufactured poultry meat products in El-Bohira governorate. Assiut Vet. Med. J., 53(112):115-124.
- Brr, A.A.H. and Mahmoud, Y.A.G. (2005). Anti-yeast effects of some plant extracts on yeasts contaminating processed poultry products in Egypt. Czech J. Food Sci., 23(1): 12-19.
- Deák, T. (2008). Handbook of food spoilage yeasts. 2nd Ed. CRC press Taylor and Francis Group, LLC.
- Edris, A.M.; Abdel-Rahman, M. and Mousa, M.M. (1992). "Microbial evaluation of some retailed cut- up chicken and poultry meat products". Benha Vet. Med. J., 3 (1): 154-166.
- Edris, A.M. (1986). mycological evaluation of frozen meat and frozen chicken. Ph. D. V. Sc. Thesis meat hygiene, Fac. Vet. Med., Alexandria Univ.
- El-Deeb, M.F.; El-Glel, H.A.; Samaha, I.A. T. (2011). Quality assurance of some poultry meat products. Alexandria Journal of Veterinary Sciences., 33(1): 153-163.
- El-Diasty, E.M.; Eman-abdeen, E. and Salem, R.M. (2013). Mycological aspects and mycotoxin residues of some chicken meat products with identification of *C. albicans* and *C. zeylanoides* by using amplified polymorphic DNA. Arab J. Biotech., 16(2):195-208.
- El-Diasty, E.M. and Salem, R.M. (2009). Incidence of lipolytic and proteolytic fungi in some milk products and their public health significance. Arab J. Biotech., 12, (1): 49-56.
- El-Gazzar, M.M.M (1995). Mycological studies on some meat products. Ph.D. Thesis Fac.

- Vet. Med. Zagazig Univ. Egypt.
- El-Khatieb, T. and Abdel-Rahman, H.A. (1989).** Mold and yeast hazard in frozen ground beef. *Assiut Vet. Med. J.*, 21(41):123-129.
- Farag R.N.E (2000).** lipolytic and proteolytic activities of some molds in some meat products. Ph.D. Thesis, Faculty of Vet. Med., Zagazig Univ., Egypt.
- Hassan, M.A. (2007).** Caprine aflatoxicosis. Ph. D. Thesis, Clinical Pathology Dept., Fac. Vet. Med., Cairo Univ., Egypt.
- Hegazi, S.M.; El-Far, F.; Edris, A.M. and Aziz, N.A. (1992).** "Studies of fungal and aflatoxins contamination of meat, meat product, and food additives". *Vet. Med. J. Giza*, 40: 31-36.
- Hussein, M.A.M. (2008).** Mycological Aspect of Fresh and Processed Meat Products with Special Respect to Proteolytic and Lipolytic Mold. Ph.D. Thesis, (Meat Hygiene) Fac. Vet. Med., Zagazig Univ., Egypt.
- IARC (1993).** IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol (56). Some naturally occurring substances: Food items and constituents, heterocyclic aromatic amines and mycotoxins, Lyon, International Agency for Research on Cancer.
- ISO (217-1-2:2008) EAST AFRICAN STANDARD.** Microbiology of food and animal feeding stuffs — Preparation of test samples, initial suspension and decimal dilutions for microbiological examination- Part 1 -3: Specific rules for the preparation of meat and meat products, 2008.
- Jay, J.M. (1979).** Modern food microbiology. 2nd Ed. D. Lvan No strand Company.
- Koburger, J.A. and Jacger, K.E. (1987).** "Specific and sensitive plate assay for bacterial lipases." *Appl. Environ. Microbiol.*, 53:211-215.
- Koburger, J.A. (1972).** Fungi in food III. The enumeration of lipolytic and proteolytic organisms *J. Milk Food. Technol.*, 35:117.
- Kocic'-Tanackov, S.D.; Dimic', G.R. and Karali c', D. (2007).** Contamination of spices with moulds potential producers of sterigmatocystine. *APTEFF*, 38: 1-190.
- Korashy- Eman and Wahbba -Nahed, M. (2008).** lipolytic and Proteolytic activities of some fungi isolated from raw camel's milk. *Assiut Vet. Med. J.*, 54: 118.
- Kruger Van Rij, N.J.W., (1984).** The yeasts: A taxonomic study. 3rd Ed. Amsterdam, Elsevier.
- Nasser, L.A. (2002).** "Mycological status of imported canned fish consumed in Saudia Arabia with special reference to proteolytic activity". *Assuit Vet. Med. J.*, 47:125-135.
- O'reilly, T. and Day, D.F. (1983).** "Effects of culture conditions on protease production by *Aeromonas hydrophila*." *Appl. Environ. Micro.*, 45: 1132.
- Ouf, J.M.; Nagwa, I.M.K. and Shabana, E.S.E. (2010).** Incidence of proteolytic and lipolytic moulds and yeasts in some ready to eat meat products. *Assuit Vet. Med. J.*, 56 (126):132-143.
- Pitt, J.I. and Hoching, A.D. (2009).** Fungi and Food spoilage. 3rd Ed. Published by Springer Dordrecht Heidelberg London New York.
- Rwaih-Wadee, E. (2010).** Mycological aspects of some chicken meat products. M.V. Sc. Thesis, Meat Hygiene, Fac. Vet. Med. Zagazig University.
- Samaha, H.A.M. (2013).** Frequency Rates of Fungal Contaminants in Imported Meats from Alexandrian Retail Markets. *Life Sci J.*, 10(4):158-165.
- Schmidt-Lorenz, W. and Gutschmidt, J. (1969).** Mikrobielle und sensorische Veränderungen gefrorener Brathähnchen und Poularden bei Lagerung im Temperaturbereich von -2.5°C bis -10°C . *Fleischwirtschaft*, 49: 1033-1041.
- Seham- Ismail, A.; Amal- Shehata, A. and El-diasty, E.M. (2013).** Microbiological quality of some meat products in local markets with special reference to mycotoxins. *Global Veterinaria*, 10 (5): 577-584.
- Shaltout, F.A. (2002).** Microbiological aspect of Semi-cooked chicken meat products. *Benha Vet. Med. J.*, 13(2):15-26.
- Shaltout, F.A. (1996).** "Mycological and My-

- cotoxicological profile of some meat products". Ph. D. Thesis Faculty Vet. Med., Moshtohor (Zagazig Univ., Benha Branch).
- Sharaf, E.M. and Sabra, S.M. (2012).** Microbiological loads for some types of cooked chicken meat products at Al-Taif governorate, KSA. World Applied Sciences Journal, 17(5): 593-597.
- Shawish, R.R.M. (2011).** Microbial evaluation of some retailed cut-up chicken and poultry meat products. M.V. Sc. Thesis, Meat Hygiene, Fac. Vet. Med. Menoufia Univ.
- Soliman, Z.I. and Shalaby, A.M. (2008).** Occurrence of lipolytic and proteolytic fungi in locally smoked fish. Assiut Vet. Med. J., 54 (117): 78-88.
- Subash, G.; Periasamy, A. and Azariah, H. (2005).** Extracellular enzymatic activity profiles in fungi isolated from oil-reach environments. J. Mycoscience, 46 (2):119-126.
- Viljoena, B.C.; Geornarash, I.; Lamprecht, A. and Von Holyb, A. (1998).** Yeast populations associated with processed poultry. Food Microbiology, 15(1):113-117.
- Welthagen, J.J. and Viljoen, B.C. (1999).** "The isolation and identification of yeasts obtained during the manufacture and ripening of Cheddar cheese." Food Microbiol., 16:63.
- Wong, C.F.; Yan, S.W. and Chan, K.W. (2008).** Report of two cases of tumour-like blocked pulmonary cavities with liquid content infected with Aspergilli, a rare form of pulmonary aspergillosis. Respiratory (13), 306-308.
- Yousif, E.I.; Ashoush, I.S.; Donia, A.A. and Hala- Goma, K.A. (2013).** Critical control points for preparing chicken meals in a hospital kitchen. Annals of Agricultural Science, 58(2): 203-211.
- Zhang, L.; Davis, M.A. and Conner, D.E. (2001).** Poultry- borne pathogens: plant considerations. Poultry meat processing chapter (9) ISBN.

مدي تواجد الفطريات المحللة للبروتين والدهون في بعض منتجات لحوم الدجاج وأهميتها الصحية

***فهم عزيز الدين شلتوت و **إيمان محمود الدياسطي و ***محمد سامي محمد**

***كلية الطب البيطري بمشتهر - جامعة بنها**

**** قسم الفطريات بمعهد بحوث صحة الحيوان بالدقي**

***** مصانع اجيبب فود**

الملخص العربي

تم في هذه الدراسة تجميع عدد ١٠٠ عينة بطريقة عشوائية من منتجات لحوم الدواجن بواقع ٢٥ عينة من كل من فيلية الدجاج و لانشون الدجاج والبانيه ولحم الدجاج المفروم من الأسواق المختلفة . وقد تم فحص هذه العينات لتقييمها من حيث تعرضها للتلوث بالفطريات المختلفة . وقد وجد أن متوسط العدد الكلي للأعفان بتلك العينات هو كالتالي (٩ , ١٠ x ٢ ± ٨ , ١٠ x ٣ , ٣ , ١٠ x ٢ , ٠ ± ١٠ x ٢ , ٨ ; ١٠ x ١ , ٤ ± ١٠ x ١ , ٩ و ١٠ x ٣ , ٤ ± ١٠ x ١ , ٩) مستعمرة / جرام لكل من عينات فيلية الدجاج و لانشون الدجاج والبانيه ولحم الدجاج المفروم على الترتيب . أما بالنسبة للعد الكلي للخمائر فقد كان ١٠ x ٣ , ٧ ± ١٠ x ٥ , ٧ و ١٠ x ٣ , ٧ ± ١٠ x ٣ , ٣ ; ١٠ x ٣ , ٣ ± ١٠ x ٢ , ٠ ; ١٠ x ٢ , ١ ± ١٠ x ١ , ١ و ١٠ x ٣ , ٣ ± ١٠ x ٤ , ٨ ± ١٠ x ٩ , ٨ مستعمرة / جرام) . وقد كانت أكثر الأعفان التي تم عزلها من العينات هي فطر الأسبرجيليس يليه البنسيليوم ثم الجيوتركيم ، الفيوزاريوم ، الكلاسيبوريم ، الميوكور ، الأيوبنسيليوم ، الأسكوبيلولاريوبسيس وأخيرا الأكرمنيوم . وبالنسبة للخمائر الأكثر عزلا من عينات منتجات لحوم الدواجن فقد كانت الكانديدا (٥ , ٣٢ %) يتبعها الرودوتيرالا (١ , ٢٢ %) ، السكارومييسيس (٢ , ١٨ %) ، التيريويسيس (٥ , ١٥ %) وأخيرا خمائر الكريبتوكوكس (٧ , ١١ %) . تم تقييم المعزولات من الأعفان والخمائر محل الدراسة لنشاطها الأنزيمي المحلل للبروتين والدهون على البيئات الخاصة بهذا الغرض . وقد تم مناقشة الأهمية الاقتصادية والصحية للفطريات التي تم عزلها الى جانب مناقشة الاحتياطات الصحية لتجنب مشاكل تلك الفطريات.