

An overview on lactoferrin as a natural antimicrobial in food industry Taghreed, H. Abbas* and Shereen, A. Yassin**

*Reference Lab for Safety Analysis of Food of Animal Origin, Food Hygiene Department; ** Food Hygiene Department, Kafr-Elsheikh Branch
Animal Health Research Institute, Agricultural Research Center (ARC), Giza, Egypt

Review Article

Corresponding author:

Taghreed Hamdy Abbas

taghreed7amdy@gmail.com

Received in 4/8/2024

Accepted in 2/9/2024

Abstract

Milk proteins have equitable composition of amino acids assuring a fundamental functional product, and they also enhance the texture and sensory qualities of various dairy products. Among them is lactoferrin (LF), which is effective against a wide range of bacteria and has both bacteriostatic and bactericidal activity. The major iron-binding protein in milk is lactoferrin (LF), which is considered as a multifunctional protein that exists in human as well as animal milk that helps adjusting the amount of free iron in body fluids; this makes LF bacteriostatic and healthy. Lactoferrin (LF) is a protein belonging to the transferrin family that is found in milk and colostrum, as well as body fluids such as saliva, tears, spleen, stomach mucosa, lymph nodes, skin, and even white blood cells. As lactoferrin binds iron, so it is depriving growth of microorganisms such as *Bacillus stearothermophilus*, *E. coli*, *Bacillus subtilis* and *Shigella* from its necessary building blocks, it also appears to serve as a protective factor against infections from bacteria, viruses, and fungi. In addition to prolonging shelf life and being used in infant formulas, it acts as a natural antimicrobial agent for the bio-preservation of dairy, seafood, meat, beverages and bakery goods, and protects against serious diseases such as hepatitis, cancer, foodborne illnesses, and respiratory infections, also helps to ensure safety and improve health. Briefly, the current review aimed to shed light on the safety and importance context of lactoferrin in food industry field as well as the benefits for human health.

Keywords: *Lactoferrin, food, preservation, shelf-life, health benefits.*

Introduction

Food antimicrobials are a variety of compounds that are employed as preservatives and reduce the microbial deterioration of food. They might impede the growth of microbes in food by disrupting its vital functions. Additionally, Antimicrobials are commonly used as preservatives in a wide range of food products. In order to improve food quality, prolong shelf life, and guarantee the final product is free of microbiological contamination **Bouarab et al., (2019)**. Consumers find it bothersome when chemical preservatives are used in processed

meals, and their demand for natural foods is growing daily **Perez et al. (2012)**. People are drawn to natural additives because they are conscious of their nutrition **Asghar et al. (2011)**. Lactoferrin is considered as a promising when implemented as an antimicrobial for production of different food products medicines, biologically active additives, and cosmetic preparations. Lactoferrin acquires plentiful health benefits regarding antifungal, antibacterial, antiparasitic, antiviral, and antitumor effects, it affirms bone growth, safeguards the intestinal epithelium, and promotes recovery of

immune system functions in animals. Lactoferrin is incorporated in hepatitis- C treatment, and also enhance health condition of patients with Alzheimer's disease **Hao et al. (2019)**. The average level of natural LF in cow's milk is around 0.2 g/L, but its maximum content can reach as high as 50 to 100 g/L in cow milk while, in human milk, it reach 2–4 g/L, and 6 to 8 g/L in human colostrum **Kai et al. (2002)**. Moreover, LF plays a defensive function against infections of gastrointestinal system and participates to secretory immune function systems **Mayeur et al. (2016)**. Experimentally, it was proved that lactoferrin protects from gastritis caused by *Helicobacter pylori*, reduces virulence during infection with enterotoxigenic *E. coli*, and sustains the integrity of intestinal mucosa **Brodsky et al. (2013)**. It was granted that LF has bacteriostatic characteristics toward gram-negative pathogens that needs extreme iron as coliforms, and it still behaves in opposition to microorganisms such as *Bacillus* species, *S. aureus*, and *L. monocytogenes* due to the interaction between LF and lipo-polysaccharides of microorganisms exposed sheet causing change in permeability **Niaz et al. (2019)**. **Binyameen et al. (2021)** revealed that consuming LF fortified milk is linked to higher iron levels in the body with enhancing iron absorption through the intestines of breast-feeding infants. Furthermore, purified LF is applied in canned food, pharmaceutical preparations, sports nutrition, oral care products and cosmetics. The inclusion of products with LF offers numerous benefits, such as the preventing the infection development, enhancement of immune function, as well as anti-inflammatory and antioxidant effects. General uses of bovine LF in food production including fermented milk and milk products, dietary iron supplements, infant and cosmetics formulas, beverages, chewing gums, immune-enhancing nutraceuticals, feed and pet care supplements, and inhibits bacterial adhesion on abiotic surfaces. Lactoferrin plays a crucial role in safeguarding the mucosal barrier by preventing viral and microbial assaults. Moreover, it has been found that LF effectively combats the formation of biofilms, and when administered orally, it has successfully treated respiratory and oral infections. Lactoferrin has been certified as GRAS prestige by the **Niaz et al.**

(2019), it has been employed as health-promoting additives in commercially available food items, and it has also gained approval as a food ingredient in products by the European Food Safety Authority **Tomita et al. (2009)**.

History of Lactoferrin:

In 1939, LF was firstly discovered by **Sorensen and Sorensen, (1940)**, as a red protein in whey, it was extracted and refined from milk of human and bovine in 1960. Lactoferrin was related to serum transferrin structure, LF is categorized as a member of the Transferrin family as mentioned by **Lambert et al. (2005)**.

Functional Properties of LF:

The particular structure of LF has been given a lot of nutritional and medical value, because it transfers iron and eliminates free radicals from body fluids; LF is essentially regarded as safe and efficient component for delivering iron to individuals who are iron deficient. **Gupta et al. (2016)** and **Lopez et al. (2016)**.

Anti-microbial activity of LF:

The iron-free form of LF, or apo-LF, is thought to contribute in iron deficiency, which is necessary for the continuation of essential microbial processes and, ultimately, a slower rate of bacterial growth **Law and Reiter (1977)**. Laterally, numerous investigations have shown that the growth of various bacterial strains were inhibited by holo-LF, the iron saturated form of lactoferrin **Ellison 3rd and Giehl (1991)**; **Ochoa and Cleary (2009)**; **Ostan et al. (2017)**. Also, it has been observed that LF exhibits antiviral properties during the initial stages of infection, this can occur through the prevention of host cell infection or in the event that host cells have already been infected, by preventing the growth of specific viruses. **Su-perti et al. (1997)**; **Marchetti et al. (1999)**; **Ikeda et al. (2000)**; **Beljaars et al. (2004)**. Lactoferrin could target a wide range of fungal species, as it can effectively eliminating *Candida albicans* and *Candida kruseibdue* due to its potent iron (Fe³⁺) content **Al-Sheikh (2009)**.

Food applications:**Bio- preservation:**

In the dairy industry, it was convinced that lactoferrin improves the sensory characteristics of Kariesh cheese that has been experimentally contaminated with certain foodborne pathogens such as *E. coli* O157:H7, *S. aureus*, *B. cereus* and *L. monocytogenes*, in addition, to extending its shelf-life at refrigerated storage **Ombarak et al. (2019)**. **Azhar et al. (2022)** achieved that LF exhibiting variable effects on the viability of *E. coli* compared to *S. aureus*, as well as *E. coli* count in Kariesh and Dominate cheese. In this regard, using of 2% of LF can inhibit *S. aureus* viability in Kariesh cheese significantly. **Da Silva et al. (2012)** concluded that LF prevent *S. aureus* population to increase in the processed cheese. **Al Habty and Ali (2022)** reported that LF favored a convenient food preservative in yoghurt due to powerful antibacterial activity against multi-drug resistant (MDR) *S. aureus* and its perfect sensorial properties, as well as LF is considered as an assuring tool for controlling growth of *E.coli* and *S. aureus* in milk and Kariesh cheese and investigated that the LF application improved the overall sensory qualities and extended the shelf-life of Kariesh cheese dramatically. In a study obtained by **Shashikumar and Puranik (2011)** and **Yassin and Abd Elhady (2023)** they noticed that LF included into cheese at different levels (10, 15, and 20 ppm) resulted in fundamental reduction in bacterial growth, such reduction increased as the LF concentration increased, ultimately leading to a substantial extension of the cheese shelf-life when compared to the control samples. **Taha et al. (2019)** declared that LF could become a bright method to decline *E. coli* non-O:157 (*E. coli* O1 and O26) growth in pasteurized milk, decrease *E.coli* non-O:157 associated illness in humans consequently.

Regarding industry of meat products, LF has an efficient role in meat preservation. **Colak et al. (2008)** pronounced that total aerobic bacteria, total psychrophilic bacteria, *pseudomonas spp.*, *coliform*, *E. coli*, yeast, and mold counts in meatballs were significantly reduced by LF treatment alone and when used in combined with nisin. The meatballs treated with this combination showed the largest reduction, extending the refrigerated shelf-life up to 10

days. Using LF at concentrations of 3% and 5% for treatment of meat samples that were contaminated with *E. coli* O157:H7 led to a reduction of 2 logs by the 9th day of storage at 10°C **Soyer et al. (2020)**. The impact of LF treatment was positively assessed the shelf-life of chilled chicken breast during storage at 4°C. Significance difference (P<0.05) was noticed between the scores of sensory traits regarding chilling times and different LF concentrations, but no significant difference (P>0.05) in taste scores. Supplementary, the concentration of 20 mg/g LF treated group had the best sensory scores and microbiological counts all over the refrigerating times until the 13th day of storage **Elsayed and Hussei (2022)**. **Del Olmo et al. (2012)** demonstrated that contaminated chicken breast samples with bacterial cultures, followed by treatment with LF and its derivatives (0.5 and 5 mg/g) concentration, and exposed to high hydrostatic pressure processing (HHP) at 400 MPa for 10 min at 10°C resulted in great reduction in bacterial counts, and a good quality of sensorial aspects.

Food packaging:

It is expected that further expansion of research related to the field of packaging materials with antimicrobial properties will occur in the near future, as it will create a significant improvement in preserving food safety and quality for as long as feasible **Niaz et al. (2019)**. The combination of natural antimicrobials and edible films may be a good way to address consumer concerns about food safety and act as a physical barrier and antimicrobial mediator **Montone et al. (2023)**. Lactoferrin is known for its antioxidant properties, enhancing antioxidant activity and helping in reducing the oxidative stress (ROS) **Gould (1996)**. Utilization of LF as nutritional supplement is Generally Recognized as Safe (GRAS) as mentioned by the USFDA and a narrative food component by EFSA. Additionally, LF exhibits remarkable heat resistance, making it highly effective in food preservation **Martinez – Hernandez et al. (2017)**; **Mulder et al. (2008)**. **Brown et al. (2008)** declared that LF can be applied in chitosan edible film altering *E. coli* and *L. monocytogenes* growth. Moreover, when combined with Lysozyme, LF and chitosan films considerably decline the growth of *E. Coli* O157:H7

and *L. monocytogenes* by about 3-log.

The application of an alginate-based coating activated with hydroxyapatite/lactoferrin/ quercetin complexes, has also demonstrated a high ability to inhibit the growth of spoilage microorganisms in fresh meat products. Additionally, after being stored for 15 days at 4 °C, it retards the generation of TVB-N and delays the changes in firmness and sensory attributes of both cooked and uncooked pork fillets **Montone et al. (2023)**. **Rollini et al. (2016)** produced a film contained LF and lysozyme, proving to be highly efficient in combating hydrogen sulfide-producing bacteria found in salmon fillets. A plasma coating functionalized with lactoferrin B was found to be a good tool to control deterioration caused by microorganisms like *Pseudomonas* and extending the shelf-life of cheese **Quinteri et al. (2013)**. **Barbieri et al. (2012)** developed a cellulose-based active packaging material that included lysozyme and LF, a release of antimicrobials was observed due to the synergism between the two components in experiments conducted with prevalent food contaminants. An edible coating of whey protein fortified with LF was studied by **Muhammad et al. (2023)**, and showed a great effect on the microbial growth in soft white cheese by eliminating or reducing of proteolytic, lipolytic bacteria, yeast and molds, as well as protect food from unfavorable storage conditions, which led to a prolongation of storage time as compared to other treatments coated or uncoated with any other additives. Many other applications of LF in was conducted in food and dairy industries as freeze-drying and ultrafiltration dairy industries, remarkably bioactive and pure LF, allowing LF incorporation in infant formula **Giansanti et al. (2016)**. Through conducting this study, it is expected to obtain a good effect of lactoferrin in preserving foods of animal origin, reducing microbial contamination, preserving product quality, and protecting consumer health from food-borne pathogens.

Conclusion

Lactoferrin application as biologically active protein is widely developed. Lactoferrin supplementation in food industry should be taken into consideration because it is believed to hold

promise for the supply of food products with an iron-binding component, enhancing their antioxidant and immunomodulating properties, and expanding the shelf life of the products. Additionally, the use of LF in food products can change their quality by increasing their healthy properties and extending shelf-life. Other advantage of LF: as an iron-containing protein, it can be used to treat anemia, primarily in expectant mothers and their newborn children. Allow for higher milk production rates to meet the needs of residents in developing nations. Establish appropriate and affordable industrial and on-farm technology to extract high-value LF protein from milk and use it for bio-preserving fresh, cut fruits, vegetables, meat, and milk products to extend shelf life, prevent disease, and improve public health.

Acknowledgments:

Not applicable.

Compliance with ethical standards.

Conflict of interest: the authors declares that they have no competing interest.

Ethical statement: the authors confirm that the ethical policies of the journal, as noted on journals author guidelines page, have been adhered to. No ethical approval was required as this is a review article with no original research data.

Consent to participate: Not applicable.

Consent for publication: Not applicable.

Code availability: Not applicable.

References

- Al Habty, S.H. and Ali, D.N. (2022)**. Efficiency of lactoferrin to eradicate multidrug resistant *Staphylococcus aureus* isolated from some dairy products. *Anim. Vet. Sci.* 11(91): 35-44. Doi10.17582/journal.aavs/2023/11.1.35.44
- Al- Sheikh, H. (2009)**. Effect of lactoferrin and iron on the growth of human pathogenic *Candida* species. *Pakistan J. of Biol., Sci.*, 12 (1):91-94. Doi:10.3923/pjbs.2009.91.94.
- Asghar, A.; Anjum, F.M. and Allen, J.C. (2011)**. Utilization of dairy byproduct proteins, surfactants, and enzymes in frozen dough. *Critical reviews in food science and*

- nutrition, 51(4), 374-382. doi: 10.1080/10408391003605482.
- Azhar, M.H.; Bebawy, H.T.; Jakle, H.R. and Walaa, S.H. (2022).** Using lactoferrin as a trail to control E. coli and staph. Aureus isolated from some types of cheese. Assiut Veterinary Medical Journal, 68(174), 49-57.
- Barbiroli, A.; Bonomi, F., Capretti G, Iametti S, Manzoni, M.; Piergiovanni, L.; and Rollini, M. (2012).** Antimicrobial activity of lysozyme and lactoferrin incorporated in cellulose-based food packaging. Food Control, 26(2): 387-392. doi: 10.1016/j.foodcont.2012.01.046.
- Beljaars, L.; Strate, B.W.; Bakker, H.I.; Rker-Smit, C.; Weigmans, F.C. and Harmsen, M.C. (2004).** Inhibition of cytomegalovirus infection by lactoferrin in vitro and in vivo. Antiviral Res., 63(3):197-208. Doi: 10.1016/j.antiviral.2004.05.002.
- Binyameen, B.; Rahim, M.A.; Khalid, W.; Nasir, N.; Ilyas, S.; Afzal, F. and Israr, H. (2021).** A Comprehensive Review on Functional Role and Additive Application of Lactoferrin. Acta Scientific Nutritional Health (ISSN: 2582-1423), 5(10): 9-16.
- Bouarab, C.L.; Degraeve, P.;Ferahout, H.;Bouajila, J. and Oulahal, N. (2019).** Plant antimicrobial polyphenols as potential natural food preservatives. Journal of the Science of Food and Agriculture, 99(4), 1457-1474. Doi:10.1002/jsfa.9357.
- Brodsky, I.B.; Bondarenko, V.M.; Tomashevskaya, N.N.; Sadchikova, E.R. and Goldman, I.L. (2013).** Antimicrobial, immunomodulating and prebiotic properties of lactoferrin. Bulletin of the Orenburg Scientific Center of the Ural Branch of the Russian Academy of Sciences, 4, 3.
- Brown, C.A.; Wang, B. and Oh, J.H. (2008).** Antimicrobial activity of lactoferrin against foodborne pathogenic bacteria incorporated into edible chitosan film. J. of Food Prot., 71 (2), 319-324. Doi: 10.4315/0362-028x-71. 2. 319.
- Colak, H.; Hampikyan, H.; Bingol, E.B. and Aksu, H. (2008).** The effect of nisin and bovine lactoferrin on the microbiological quality of Turkish-style meatball (TekirdağKöfte). Journal of Food Safety, 28(3), 355-375. doi:10.1111/j.1745-4565. 2008.00105.x.
- Da Silva, A.; Honjoya, E.R.; Cardoso, S.C; Batista-De Souza, C.H; De Rezende C.M.; Walter-De Santana, E.H. and Aragon-Alegro L.C. (2012).** Antimicrobial action of lactoferrin on Staphylococcus aureus inoculated in Minas frescal cheese. Archivoslatinoamericanos de nutricion, 62(1), 68-72.
- Del Olmo, A.; Calzada, J. and Nuñez, M. (2012).** Effect of lactoferrin and its derivatives against gram-positive bacteria in vitro and, combined with high pressure, in chicken breast fillets. Meat science, 90(1), 71-76. doi:101016/j.meatsci.2011.06.003.
- Ellison 3rd, R.T. and Giehl, T.J. (1991).** Killing of gram- negative bacteria by lactoferrin and lysozyme. J Of Clin. Investi., 88(4): 1080-91. doi:10.1172/JCI11407.
- Elsayed, H. and Hussei, D.E. (2022).** Impact of Different Concentrations of Lactoferrin on the Shelf Life of the Chilled Chicken Breast. Taiwanese Journal of Agricultural Chemistry and Food Science, 60 (3).doi:10.6578/TJACFS.202209.60(3).0005.
- Giansanti, F.; Panella, G.; Leboffe, L. and Antonini, G. (2016).** Lactoferrin from milk: Nutraceutical and pharmacological properties. Pharmaceuticals, 9(4): 61. doi:10.3390/ph9040061.
- Gould, G.W. (1996).** industry perspectives on the use of natural antimicrobial and inhibitors for food applications. J. Food Prot., 59 (13): 82-86. Doi: 10.4315/0362-028X-59.13.82.
- Gupta, P.M.; Perrine, C.G.; Mei, Z. and Scanlon, K.S. (2016).** Iron, anemia, and iron deficiency aneamia among young children in the United States, Nutrients, 8(6):330.

Doi:10.3390/nu8060330.

Hao, L.; Shan, Q.; Wei, J.; Ma, F. and Sun, P. (2019). Lactoferrin: major physiological functions and applications. *Current Protein and Pept. Sci.*, 20(2), 139-144. doi: 10.2174/1389203719666180514150921.

Ikeda, M.; Nozaki, A.; Sugiyama, K.; Tanaka, T.; Naganuma, A. and Tanaka, K. (2000). Characterization of antiviral activity of lactoferrin against hepatitis C virus infection in human cultured cells. *Virus Research*, 66(1): 51-63.

Kai, K.; Komine, Y.; Komine, K.I.; Asai, K.I.; Kuroishi, T.; Kozutsumi, T. and Kumagai, K. (2002). Effects of bovine lactoferrin by the intramammary infusion in cows with staphylococcal mastitis during the early non-lactating period. *J of Vet Med Sci.*, 64(10), 873-878. doi:10.1292/jvms.64.873.

Lambert, L.A.; Perri, H. and Meehan, T.J. (2005). Evolution of duplications in the transferrin family of proteins. *Comparative Biochemistry and physiology Part B: Biochemistry and Molecular Biology*, 140(1):11-25. doi: 10.1016/j.cbpc.2004.09.012.

Law, B.A.; and Reiter, B. (1977). The isolation and bacteriostatic properties of lactoferrin from bovine milk whey. *J. of Dairy Res.*,44(03): 595-599. Doi: 10.1017/s0022029900020550.

Lopez, A.; Cacoub, P.; Macdougall, I.C. and Peyrin, B.L. (2016). Iron deficiency, anaemia. *The Lancet*, 387(10021), 907-916. Doi:10.1016/S0140-6736(15)60865-0.

Marchetti, M.; Superti, F.A.; Ammendolia, M.G. Rossi, V.P. and Seganti, L. (1999). Inhibition of poliovirus type 1 infection by iron- manganese and zinc saturated lactoferrin. *Med.Microbiol. And Immunol.* 187(4): 199-204. Doi:10.1007/s004300050093.

Martinez – Hernandez, G.B.; Amodia, M.L.; Valera- de Chiara, M.L.; Colelli, G. (2017). Microbial inactivations with hydro-

lysed lactoferrin and other natural antimicrobials in fresh- cut fennel. *LWT.*84,353-358. Doi:10.1016/j.lwt.2017.05.079.

Mayeur, S.; Spahis, S.; Pouliot, Y. and Levy, E. (2016). Lactoferrin, a pleiotropic protein in health and disease. *Antioxidand Redox Signa.*, 24(14), 813-836. Doi:10.1089/ars.2015.6458.

Montone, A.M.; Malvano, F.; Taiano, R.; Capparelli, R.; Capuano, F. and Albanese, D. (2023). Alginate Coating Charged by Hydroxyapatite Complexes with Lactoferrin and Quercetin Enhances the Pork Meat Shelf Life. *Foods*, 12(3), 553. doi:10.3390/foods12030553.

Muhammad, J.M.; Somaya, K.B. and Muhammad, Y.A. (2023). Evaluate the effect of using a whey protein membrane loaded with zinc nanoparticles, lactoferrin and neptomycin on microbial growth in laboratory-made soft white cheese. *Med. and Clin. Res.*, 8 (5), 1-8. www.medclinres.org

Mulder, A.M.; Connellan, P.A.; Oliver, C.J.; Morris, C.A. and Stevenson, L.M. (2008). Bovine lactoferrin supplementation supports immune and antioxidant status in healthy human males. *Nutr. Res.*, 28(9): 583-589. doi: 10.1016/j.nutres.2008.05.007.

Niaz, B.; Saeed, F, Ahmed, A.; Imran, M.; Maan, A.A.; Khan, M.K. and Suleria, H.A. (2019). Lactoferrin (LF): A natural antimicrobial protein. *Intern. J of Food Properties*, 22(1), 1626-1641. 0Doi: 10.1080/10942912.2019.1666137.

Ochoa, T.J. and Cleary, T.G. (2009). Effect of lactoferrin on enteric pathogens. *Biochimie.*, 91(1):30-34. doi:10.1016/j. biochi.2008.04.006.

Omarak, R.A.; Saad, M.A. and Elbagory, A.R.M. (2019). Biopreservative Effect of Lactoferrin Against Foodborne Pathogens Inoculated in Egyptian Soft Cheese" *Karish Cheese". Alexandria Journal for Veterinary Sciences*, 63(2): 97. Doi:10.5455/ajvs.76015.

- Ostan, N.K.; Yu, R.H.; Ng, D.; Lai, C.C.; Pogoutse, A.K. and Sarpe, V. (2017).** Lactoferrin binding protein 8-a bi- functional bacterial receptor protein. *PLoS Pathogens*, 13(3): 1006244. doi: 10.1371/journal. ppat. 1006244.
- Perez Espitia, P.J.; de Fátima Ferreira Soares, N.; dos Reis Coimbra, J.S.; de Andrade, N.J.; Souza Cruz, R. and Alves Medeiros, E.A. (2012).** Bioactive peptides: Synthesis properties, and applications in the packaging and preservation of food. *Comprehensive Reviews in food Sci. and Food Safety*, 11(2):187-204. Doi: 10.1111/j.1541-4337. 2011.00179. x.
- Quinteri, L.; Pistillo, P.B.; Caputo, L.; Favia, P. and Baruzzi, F. (2013).** Bovine lactoferrin and lactoferricin on plasma-deposited coating against spoilage *Pseudomonas* spp. *Innovative Food Science and Emerging Technologies*. 20(4): 215–222.doi: 10.1016/j.ifset.2013.04.013.
- Rollini, M.; Nielsen, T.; Musatti, A.; Limbo, S.; Piergiovanni, L.; Hernandez Munoz, P. and Gavara, R. (2016).** Antimicrobial performance of two different packaging materials on the microbiological quality of fresh salmon. *Coatings*, 6(1): 6. doi:10.3390/coatings6010006.
- Shashikumar, C.S. and Puranik, D.B. (2011).** Study on use of lactoferrin for the biopreservation of paneer. *Tropical Agricultural Res.*23(1): 70-76
- Sorensen, M. and Sorensen, S. (1940).** The protein in whey. *Compte rendu des Travaux du Labotatoire de Carlsberg Ser. Chim*, 23 (7): 55-99.
- Soyer, F.; Keman, D.; Eroğlu, E. and Türe, H. (2020).** Synergistic antimicrobial effects of activated lactoferrin and rosemary extract in vitro and potential application in meat storage. *Journal of Food Sci. and Technol.*, 57(12): 4395-4403. doi:10.1007/s13197-020-04476-5.
- Superti, F.; Ammendolia, M.G.; Valenti, P. and Seganti, L. (1997).** Antrotaviral activity of milk proteins: Lactoferrin prevents rotavirus infection in the enterocyte- like cell line HT -29. *Medical Microbiol. And Immunol.*, 18(2):83-91. Doi:10.1007/s004300050049.
- Taha, N.; El Barbary, H.A.; Ibrahim, E.M.; Mohammed, H.A. and Wahba, N.M.M. (2019).** Application of lactoferrin as a trial to control *E. coli* O1and O26 in pasteurized milk. *Benha Veterinary Medical Journal*, 36 (2), 360-366.
- Tomita, M.; Wakabayashi, H.; Shin, K.; Yamauchi, K.; Yaeshima, T. and Iwatsuki, K. (2009).** Twenty-five years of research on bovine lactoferrin applications. *Bioc- himie*, 91(1): 52-57.doi: 10.1016/j. biochi. 2008.05.021.
- Yassin, S.A. and Abd Elhady, H.A. (2023).** Utilization of lactoferrin to inhibit *E. coli* and *S. aureus* isolates from milk and kariesh cheese. *Iraqi Journal of Veterinary Scienc- es*, 37(4), 853-863.doi:10.33899/IJVS. 2023. 137384.2677.